

## References for misoprostol article

### References

1. Child TJ, Thomas J, Rees M, MacKenzie IZ. Morbidity of first trimester aspiration termination and the seniority of the surgeon. *Hum Reprod* 2001; **16**: 875-78
2. Joint study of the Royal College of General Practitioners and the Royal College of Obstetricians and Gynaecologists. Induction abortion operations and their early sequelae. *J R Coll Gen Pract* 1985; **35**: 175-80
3. Wamwana EB, Ndavi PM, Gichangi PB, et al. Socio-demographic characteristics of patients admitted with gynaecological emergency conditions at the provincial general hospital, Kakamega, Kenya. *East Afr Med J* 2006; **83**: 659-65
4. Child TJ, Thomas J, Rees M, MacKenzie IZ. A comparative study of surgical and medical procedures: 932 pregnancy terminations up to 63 days gestation. *Hum Reprod* 2001; **16**: 67-71
5. Bugalho A, Faundes A, Jamisse L, et al. Evaluation of the effectiveness of vaginal misoprostol to induce first trimester abortion. *Contraception* 1996; **53**: 243-46
6. Carbonell Esteve JL, Varela L, Velazco A, et al. Vaginal misoprostol for late first trimester abortion. *Contraception* 1998; **57**: 329-33
7. Carbonell Esteve JLL, Varela L, Velazco A, et al. Early abortion with 800 µg of misoprostol by the vaginal route. *Contraception* 1999; **59**: 219-25
8. Carbonell JLL, Rodríguez J, Aragón S, et al. Vaginal misoprostol 1000 µg for early abortion. *Contraception* 2001; **63**: 131-36
9. Jain JK, Meckstroth KR, Mishell Jr DR. Early pregnancy termination with intravaginally administered sodium chloride solution-moistened misoprostol tablets: historical comparison with mifepristone and oral misoprostol. *Am J Obstet Gynecol* 1999; **181**: 1386-91
10. Ngai SW, Tang OS, Chan YM, Ho PC. Vaginal misoprostol alone for medical abortion up to 9 weeks of gestation: efficacy and acceptability. *Hum Reprod* 2000; **15**: 1159-62
11. Jain JK, Meckstroth KR, Park M, Mishell Jr DR. A comparison of tamoxifen and misoprostol to misoprostol alone for early pregnancy termination. *Contraception* 2000; **60**: 353-56
12. Bugalho A, Mocumbi S, Faundes A, David E. Termination of pregnancies of <6 weeks gestation with a single dose of 800 µg of vaginal misoprostol. *Contraception* 2000; **61**: 47-50
13. Jain JK, Dutton C, Harwood B, et al. A prospective randomized, double-blinded, placebo-controlled trial comparing mifepristone and vaginal misoprostol to vaginal misoprostol alone for elective termination of early pregnancy. *Hum Reprod* 2002; **17**: 1477-82
14. Zikopoulos KA, Papanikolaou EG, Kalantaridou SN, et al. Early pregnancy termination with vaginal misoprostol before and after 42 days gestation. *Hum Reprod* 2002; **17**: 3079-83
15. Jain JK, Harwood B, Meckstroth KR, Mishell DR. Early pregnancy termination with vaginal misoprostol combined with loperamide and acetaminophen prophylaxis. *Contraception* 2001; **63**: 217-21
16. Carbonell JLL, Rodríguez J, Velazco A, et al. Oral and vaginal misoprostol 800 µg every 8 h for early abortion. *Contraception* 2003; **67**: 457-62
17. Salakos N, Kountouris A, Botsis D, et al. First-trimester pregnancy termination with 800 µg of vaginal misoprostol every 12h. *Eur J Contracep Reprod Health Care* 2005; **10**: 249-54
18. Blanchard K, Shochet T, Coyaji K, et al. Misoprostol alone for early abortion: an evaluation of seven potential regimens. *Contraception* 2005; **72**: 91-97
19. von Hertzen H, Piaggio G, Nguyen Thi MH, et al. on behalf of the WHO Research Group on Postovulatory Methods of Fertility Regulation. Efficacy of two intervals and two routes of administration of misoprostol for termination of early pregnancy: a randomised controlled equivalence trial. *Lancet* 2007; **369**: 1938-46
20. Cheung W, Tang OS, Lee SW, Ho PC. Pilot study on the use of sublingual misoprostol in termination of pregnancy up to 7 weeks gestation. *Contraception* 2003; **68**: 97-99